Author Index

Ahuja, S
——, Thompson, G. and Smith, J.
Trace/ultratrace analyses of unstable compounds. Investigations on hydrobenzene and azobenzene 443
Akapo, S. O.
——, Furst, A., Khong, T. M. and Simpson, C. F.
Chromatographic evaluation of oligomeric C_8 reversed phases for the use in high-performance liquid chromatography 283
Apps, P. J.
—— and Pretorius, V.
Comparison between the conditions for solute focussing by the static and dynamic solvent effects under ideal conditions 81
Asano, T., see Takeuchi, T. 297
Audunsson, A., see Nilvé, G. 151
Bartle, K. D.
——, Boddington, T., Clifford, A. A., and Shilstone, G. F.
Effect of pressure drop across the column on average densities and capacity factors in supercritical fluid chromatography 347
Bertsch, W., see Green, L. S. 113
Betz, W. R.
—— and Supina, W. R.
Determination of the gas chromatographic performance characteristics of several graphitized carbon blacks 105
Bighi, C., see Pietrogrande, M. C. 407
Boddington, T., see Bartle, K. D. 347
Boesten, W. H. J., see Duchateau, A. 263
Borea, P. A., see Pietrogrande, M. C. 407
Brinkman, U. A. Th., see Lipschitz, C. 321
Bruin, G. J. M.
——, Chang, J. P., Kuhlman, R. H., Zegers, K., Kraak, J. C. and Poppe, H.
Capillary zone electrophoreic separations of proteins in polyethylene glycol-modified capillaries 429
Chang, J. P., see Bruin, G. J. M. 429
Clifford, A. A.
Theory of open-tube chromatography: an exact proof of Golay's equations 61
——, see Bartle, K. D. 347
Crombach, M., see Duchateau, A. 263
Damme, G., Van, see Verzele, M. 335
De Jong, G. J., see Lipschitz, C. 321
Dimov, N.
——, and Mekenyan, Ov.
Quantitative relationships between the structure of alkylbenzenes and their gas chromatographic retention on stationary phases with different polarity 227
Dondi, F., see Pietrogrande, M. C. 407
Drawert, F., see Nitz, S. 173
Duchateau, A.
——, Crombach, M., Kamphuis, J., Boesten, W. H. J., Schoemaker, H. E. and Meijer, E. M.
Determination of the enantiomers of α-H-α-amino acids, α-alkyl-α-amino acids and the corresponding acid amides by high-performance liquid chromatography 263
Engewald, W., see Maurer, T. 245
Evans, M. B.
—— and Haken, J. K.
Dispersion and selectivity indices in gas chromatography. Part II. Studies of homologous carbonyl and carboxyl compounds 217
Farkaš, P.,
——, Soják, L., Kováč, M. and Janák, J.
Interface adsorption and reproducibility of retention indices in glass capillary columns with dimethylpolysiloxane stationary phases cross-linked by γ-irradiation 251
Feltl, L., see Kohoutová, A. 139
Fernández-Díaz, M., see Garciá-Raso, A. 205
Finney, R. W.
—— and Read, F.
Laser microprobe mass spectrometry of selected compounds directly from normal phase high-performance thin-layer plates 389
Frei, R. W., see Lipschitz, C. 321
Furst, A., see Akapo, S. O. 283
García-Raso, A.
——, Fernández-Díaz, M. Páez, M. I., Sanz, J. and Martínez-Castro, I.
Gas chromatographic retention of cabrohydrate trimethylsilyl ethers. III. Ketohexoses 205
Gehrke, C. W.
—— and Guo, C. K.
Ribonucleoside analysis by reversed-phase high-performance liquid chromatography 3
Green, L. S.
—— and Bertsch, W.
Charatceristics of glass based packings as a support in chromatography 113

Grob, K.
——, Lanfranchi, M. and Mariani, C.
Determination of free and esterified sterols and wax esters in oils and fats by coupled liquid chromatography–gas chromatography 397
Grushka, E.
—— and McCormick, R. M.
Zone boradening due to sample injection in capillary zone electrophoresis 421
Guha, O. K.
On some nomenclature in chromatography 448
Guo, C. K., see Gehrke, C. W. 3
Haken, J. K., see Evans, M. B. 217
Heybey, J., see Mohnke, M. 37
Horká, M., see Janák, K. 237
Ioffe, B. V., see Vitenberg, A. G. 55
Irth, H., see Lipschitz, C. 321
Ishii, D., see Takeuchi, T. 297
Janák, J., see Farkaš, P. 251
Janák, K.
——, Horká, M. and Tesařík, K.
Effect of repeated cross-linking of SE-54 stationary phase film on the chromatographic properties of capillary columns 237
Janeček, M.
—— and Šlais, K.
Simultaneous photometric and conductivity detection for microcolumn liquid chromatography 303
Jönsson, J. Å., see Nilvé, G. 151
Jong, G. J. de, see Lipschitz, C. 321
Kamphuis, J., see Duchateau, A. 263
Katsanos, N. A.
—— and Vassilakos, Ch.
New method for measuring obstructive factors and porosity using gas chromatographic instrumentation 123
Kersten, B. R., see Poole, S. K. 91
Khong, T. M., see Akapo, S. O. 283
Kohoutová, A.
——, Smolková-Keulemansová, E. and Feltl, L.
Use of gas chromatography for a study of crown ethers 139
Kollmannsberger, H., see Nitz, S. 173
Kováč, M., see Farkaš, P. 251
Kraak, K., see Bruin, G. J. M. 429
Kuhlman, R. H.,m see Bruin, G. J. M. 429
Kuksis, A., see Myher, J. J. 187
Lanfranchi, M., see Grob, K. 397
Lipschitz, C.
——, Irth, H., De Jong, G. J., Brinkman, U. A. Th. and Frei, R. W.
Trace enrichment of pyrimidine nucleobases, 5-fluorouracil and bromacil on a silver-loaded thiol stationary phase with on-line reversed-phase high-performance liquid chromatography 321
Lochmüller, C. H.
—— and Mink, L. P.
Adsorption isotherms on silica for methanol and 1-hexanol modifiers from supercritical carbon dioxide 357
Lyne, P. M.
—— and Philips, C. S. G.
Isomerization of propyne to propadiene. Studies by gas chromatography 145
McCormick, R. M., see Grushka, E. 421
McCune, J. E., see Pirkle, W. H. 271
Mariani, C., see Grob, K. 397
Martínez-Castro, I., see García-Raso, A. 205
Martire, D. E.
Application of a unified molecular theory to gas–liquid chromatography 71
Maurer, T.
——, Welsch, Th., and Engewald, W.
Non-linearity of the plot of log(adjusted retention time) *versus* carbon number for *n*-alkanes in series-coupled gas chromatographic columns 245
Meijer, E. M., see Duchateau, A. 263
Mekenyan, Ov., see Dimov, N. 227
Mincsovic, E., see Tyihák, E. 375
Mink, L. P., see Lochmüller, C. H. 357
Mohnke, M.
—— and Heybey, J.
Gas–solid chromatography on open-tubular columns: an isotope effect 37
Mori, S.
Aqueous size-exclusion chromatography of anionic and non-anionic water-soluble polymers on silica gel with bonded hydrophilic groups 367
Myher, J. J.
—— and Kuksis, A.
Reative gas–liquid chromatographic retention factors of trimethylsilyl ethers of diradylglycerols on polar capillary columns 187
Nilvé, G.
——, Audunsson, A. and Jönsson, J. Å.
Sample preparation by means of a supported liquid membrane for the determination of chlorophenoxyalkanoic acids 151

Nitz, S.
——, Kollmannsberger, H. and Drawert, F.
Determination of sensorial active trace compounds by multi-dimensional gas chromatography combined with different enrichment techniques 173
Noda, H., see Tsuda, T. 311
Onuska, F.
—— and Terry, K. A.
Quantitative high-resolution gas chromatography and mass spectrometry of toxaphene residues in fish 161
Páez, M. I., see García-Raso, A. 205
Phillips, C. S. G., see Lyne, P. M. 145
Pietrogrande, M. C.
——, Dondi, F., Borea, P. A. and Bighi, C.
Retention behaviour of β-carbolines in normal-phase chromatography. Silica and amino phases in high-performance and thin-layer chromatography 407
Pirkle, W. H.
—— and McCune, J. E.
Improved chiral stationary phase for the separation of the enantiomers of chiral acids as their anilide derivatives 271
Poole, C. F., see Poole, S. K. 91
Poole, S. K.
——, Kersten, B. R. and Poole, C. F.
Comparison of solvent models for characterizing stationary phase selectivity in gas chromatography 91
Poppe, H., see Bruin, G. J. M. 429
Pretorius, V., see Apps, P. J. 81
Read, F., see Finney, R. W. 389
Roberts, D. W.
——, Ruane, R. J. and Wilson, I. D.
Picolinic acid: a mobile phase additive for improved chromatography of metal-chelating heterocyclic acids and β-diketones 437
Ruane, R. J., see Roberts, D. W. 437
Sanz, J., see Garciá-Raso, A. 205
Schoemaker, H. E., see Duchateau, A. 263
Schuddink, G., see Verzele, M. 335
Shilstone, G. F., see Bartle, K. D. 347
Šlais, K., see Janeček, M. 303
Smith, J., see Ahuja, S. 443
Smolková-Keulemansová, E., see Kohoutová, A. 139
Soják, L., see Farkaš, P. 251
Supina, W. R., see Betz, W. R. 105
Székely, T. J., see Tyihák, E. 375
Takeuchi, T.
——, Asano, T. and Ishii, D.
Mass detection limits achieved with a commercially available fluorimeter in micro high-performance liquid chromatography 297
Terry, K. A., see Onuska, F. 161
Tesařík, K., see Janák, K. 237
Thompson, G., see Ahuja, S. 443
Tsuda, T.
—— and Noda, H.
Effects of the position of the laser beam focal point on a capillary flow-through cell on the signal-to-noise ratio for a fluorimetric detector in capillary column liquid chromatography 311
Tyihák, E.
——, Mincsovic, E. and Székely, T. J.
Overpressured multi-layer chromatography 375
Van Damme, G., see Verzele, M. 335
Vassilakos, Ch., see Katsanos, N. A. 123
Verzele, M.
——, Van Damme, G., Schuddink, G. and Vyncke, P.
Quantitative microscale liquid chromatography of piperine in pepper and pepper extracts 335
Vitenberg, A. G.
—— and Ioffe, B. V.
Basic equations in continuous gas extraction and their application to headspace analysis 55
Vyncke, P., see Verzele, M. 335
Welsch, Th., see Maurer, T. 245
Wilson, I. D., see Roberts, D. W. 437
Zegers, K., see Bruin, G. J. M. 429